



# Nasjonalt Senter for Gastroenterologisk Ultrasonografi

National Centre for Ultrasound in Gastroenterology  
Haukeland University Hospital, Bergen, Norway

## Ultrasound of diffuse and focal liver diseases


Odd Helge Gilja, MD, PhD

Professor

Department of Medicine

Haukeland University Hospital


Bergen, Norway





# The Ultrasound Tool-box

- Ultrasound of liver
  - B mode
  - B-Flow
  - Doppler
    - Color Doppler
    - Pulsed Doppler
  - Elastography
    - Strain imaging
    - Shear wave
  - Contrast-US (CEUS)
- US-guided liver biopsy (Menghini and Pistol)
- US-guided ablation techniques
- Sonoporation therapy






# Ultrasound of the Liver


## -What do we look for ?

- Echogenicity
- Size, capsule and form
- Any lesions?
- Liver veins
- Portal vein
- Arteria hepatica
- Intrahepatic bile ducts





# The Liver capsule





# Vena Porta



Haukeland US

09/29/10 09:20:35

ADM

MI 1.2 TIs 1.7 C1-5


GASTRO

FR 24

CHI


0-Frq 5.0  
Gn 64  
S/A 1/1  
Map F/1  
D 12.0  
DP 66

LOGIQ  
E9






# The hepatic veins






# Doppler is useful !







# US better than CT

- Small and complex cysts
- Flow evaluation
  - Portal vein trombosis
  - Budd-Chiari
- Cirrhosis evaluation
  - Global liver assessment
  - Portal HT
  - TIPS





# Segmentation of the liver



The liver's eight-segment division. Patients undergoing surgery for liver cancer will have a section(s) of their liver removed in which the tumor(s) resides. Following surgery, this section grows back.



# 6+

## A systematic examination of abdominal organs

- **Stasjon 1** Tidlig morgen før frokost. Man skanner i lengderetning med en tom mag. Landemerker: Abdominalmarkene. Systematisk gjennomgang av magen fra dypt inn til å få et drikke kan noensinne drikke.
  - **Stasjon 2** I lengderetning over magen (antrum) i sammenheng med avgående kar i gesekken.
  - **Stasjon 3** Som man noen gang undersøkes fas med vinkling og intercostalt.
  - **Stasjon 4** Lydhodet og skallskjell i intercostalt.
  - **Stasjon 5** Som lokaliserer milte og nyre fra pol til pol. Venstre flanke gjøres best når man sitter.
  - **Stasjon 6** Urinblæren er fyldt og sees.
  - **Stasjon +** For å se etter "tagg" og identifisere den retning ned til sigmoideum.
- 
- The diagram shows a grayscale ultrasound image of a human torso from the side. Six black numbers are overlaid on the image to indicate scanning stations:
- 1**: Located at the top center, just below the diaphragm.
  - 2**: Located in the upper right quadrant, just below the liver.
  - 3**: Located on the left side, just below the liver.
  - 4**: Located on the right side, just below the liver.
  - 5**: Located on the left side, below the liver and above the kidney area.
  - 6**: Located at the bottom center, just above the rectum.



# Stasjon 2 – Ve. lobe




Haukeland US / NSGU  
01/17/18 10:58:06

ADM

MI 1.2 TIs 0.6 C1-6  
Abdomen

FR 28



CHI  
Frq 4.0  
Gn 59  
0° S/A 2/1  
- Map A/1  
- D 8.0  
- DR 72  
- AO% 100

2"

-

-

-

-

-

-

X

-

-


-

8"

1 L 0.56 cm




# Stasjon 3 – Scanne for FLLs





# Sammenligne ekkogenisitet lever-nyre







# Alkoholiske leversykdommer

- Generell parenchymaffeksjon
  - Steatose
  - Hepatitter
  - Fibrose – Cirrhose
- Fokale lesjoner
  - Regenerasjonsknuter - DN
  - HCC





# Alkoholisk Hepatitt



Hvordan spørre om  
Alkoholvaner?





"Jeg drikker bare et glass om  
dagen, doktor!"






# Alcoholic Hepatitis






# Liver cirrhosis






# Ultrasound in the evaluation of cirrhotic livers

- Size of liver
- Size of left lobe and caudate lobe
- Capsule smoothness
- Ascites
- Echogenicity, homogeneity, nodularity, focal lesions
- Bile ducts and gallbladder
- Diameter of portal vein (+ splenic vein and spleen)
- Doppler measurements:
  - Color and pulsed Doppler of portal and hepatic veins
  - Doppler of hepatic artery (TX)
- Elastography, mainly right lobe
- CEUS
- US-guided biopsy and ablation procedures





# Lever Cirrhose





# Color Doppler flow in real-time



Haukeland US  
06/02/10 11:21:28

ADM

MI 0.9 Tls 1.4 C1-5  
GASTRO

FR 24




CHI  
0° Frq 4.0  
Gn 75  
S/A 1/1  
Map F/1  
D 14.0  
DR 66  
AO% 100

5°  
10°  
15°  
20°  
25°  
30°  
35°  
40°  
45°  
50°  
55°  
60°  
65°  
70°  
75°  
80°  
85°  
90°

-20  
cm/s

LOGIQ  
E9





0°

5°

10°




# Enlarged caudate lobe






# Dilated splenic vein in Portal HT





# Esophageal Varices





# Fatty liver – Dangerous !

\$1.09/US\$1 CANADA  
Vol. 13 - No. 26 June 27, 1995

# Sun

# 600 lb wife sits on husband and kills him




# Steatose

GE Healthcare  
02/03/10 10:39:37 ADM

MI 0.9 TIs 1.4 C1-5  
GASTRO

FR 24

CHI  
Frq 4.0  
0-Gn 70  
S/A 1/1  
Map F/1  
D 14.0  
DR 66  
AO% 100




9 ADM

MI 0.8 TIs 0.8 C1-5  
GASTRO


FR 25

CHI  
Frq 3.0  
0-Gn 71  
S/A 1/1  
Map F/1  
D 13.0  
DR 66  
AO% 100







# Steatosis







# Fatty liver with "focal areas of sparing"








# Fatty liver with "focal areas of sparing"





# PSC: Primary Sclerosing Cholangitis







# Longitudinal and Shear Waves

Ultrasound  
Wave

$$c_l = \sqrt{\frac{K}{\rho}}$$

$c_l \sim 1540$  m/s  
in tissue




Shear Wave

$$c_t = \sqrt{\frac{E}{3\rho}}$$

$c_t = 1-10$  m/s in tissue



# Shear Wave Elastography compared to histological findings and Liver Fibrosis



*Ultrasound Med Biol. 2011 Sep;37(9):1361-73. Epub 2011 Jul 2011 Noninvasive in vivo liver fibrosis evaluation using supersonic shear imaging: a clinical study on 113 hepatitis C virus patients. Bavu E, Gennisson JL, Couade M, Bercoff J, Mallet V, Fink M, Badel A, Vallet-Pichard*



# New Guidelines 2017

Guidelines & Recommendations

## 29 recommendations

 Thieme

### EFSUMB Guidelines and Recommendations on the Clinical Use of Liver Ultrasound Elastography, Update 2017 (Long Version)

### EFSUMB-Leitlinien und Empfehlungen zur klinischen Anwendung der Leberelastographie, Update 2017 (Langversion)

#### Authors

Christoph F. Dietrich<sup>1, 2</sup>, Jeffrey Bamber<sup>3</sup>,  
Annalisa Berzigotti<sup>4</sup>, Simona Bota<sup>5</sup>, Vito Cantisani<sup>6</sup>,  
Laurent Castera<sup>7</sup>, David Cosgrove<sup>8</sup>, Giovanna Ferraioli<sup>9</sup>,  
Mireen Friedrich-Rust<sup>10</sup>, Odd Helge Gilja<sup>11</sup>,  
Ruediger Stephan Goertz<sup>12</sup>, Thomas Karlas<sup>13</sup>, Robert de  
Knegt<sup>14</sup>, Victor de Ledinghen<sup>15</sup>, Fabio Piscaglia<sup>16</sup>,  
Bogdan Procopet<sup>17</sup>, Adrian Saftoiu<sup>18</sup>, Paul S. Sidhu<sup>19</sup>,  
Ioan Sporea<sup>20</sup>, Maja Thiele<sup>21</sup>

13 Department for Internal Medicine, Division of  
Gastroenterology and Rheumatology, University Hospital  
Leipzig, Leipzig, Germany

14 Department of Gastroenterology and Hepatology,  
Erasmus MC University Medical Center, Rotterdam, the  
Netherlands

15 Non-invasive diagnosis of liver fibrosis centre, Haut-  
Leveque hospital, Bordeaux University Hospital, Pessac,  
France

16 Unit of Internal Medicine, Dept of Medical and Surgical



# SWE in Hepatitis C and B

## RECOMMENDATION 18

2D-SWE as demonstrated with SSI can be used as a first-line assessment for the severity of liver fibrosis in patients with chronic viral hepatitis C. It performs best with regard to the ruling out of cirrhosis (LoE 1b, GoR A) [139, 158, 159]. Broad consensus (17/0/1, 94 %)

## RECOMMENDATION 24

2D-SWE as demonstrated with SSI is useful in patients with CHB to identify those with cirrhosis (LoE 3a, GoR C) [196, 197]. Broad consensus (17/0/1, 94 %)




# Ultrasound elastography

- Correlates well with histology regarding fibrosis
- Easy to perform
- Prolongs the US exam only with 2 min
- Provides valuable information to the clinician
- CT does not give data on liver stiffness
- MR elastography has low availability, is expensive and time consuming



# Detection of Focal Liver Lesions



Hausken, Gilja et al., 1999





# FLL Classification

Benign	Malignant
Hepatocellular Adenoma Focal Nodular Hyperplasia Diffuse Nodular Hyperplasia Macroregenerative Nodules Dysplastic Nodules	Hepatocellular Hepatocellular Carcinoma (Hcc) and its variants Fibrolamellar Carcinoma Hepatocholangiocarcinoma Hepatoblastoma Carcinosarcoma
Biliary Epithelium Bile Duct Cyst Biliary Duct -Adenoma Mucinous Cystic Neoplasm Peribiliary Gland Hamartoma von Meyenburg Complex Biliary Cystadenoma Biliary Papillomatosis	Biliary Epithelium Cystadenocarcinoma Cholangiocarcinoma
Vascular Cavernous Hemangioma Infantile Hemangioendothelioma	Vascular Angiosarcoma Epithelioid Hemangioendothelioma
Others Angiomyolipoma Mesenchymal Hamartoma Solitary Fibrous Tumor Inflammatory Pseudotumor	Others Primary Lymphomas Sarcomas





# Liver cysts







# Polycystic liver disease






# sonoVue®

Sulphur Hexafluoride





# Sonazoid



- Membrane is hydrogenated egg phosphatidylserine sodium (HEPSNa)
- The gas is perfluorobutane (PFB)



# New Guidelines for CEUS 2011

## The EFSUMB Guidelines and Recommendations on the Clinical Practice of Contrast Enhanced Ultrasound (CEUS): Update 2011 on non-hepatic applications

### Authors

F. Piscaglia<sup>1</sup>, C. Nolsøe<sup>2</sup>, C. F. Dietrich<sup>3</sup>, D. O. Cosgrove<sup>4</sup>, O. H. Gilja<sup>5</sup>, M. Bachmann Nielsen<sup>6</sup>, T. Albrecht<sup>7</sup>, L. Barozzi<sup>8</sup>, M. Bertolotto<sup>9</sup>, O. Catalano<sup>10</sup>, M. Clouston<sup>11</sup>, D. A. Clevert<sup>12</sup>, J. M. Correas<sup>13</sup>, M. D'Onofrio<sup>14</sup>, F. M. Drudi<sup>15</sup>, J. Eyding<sup>16</sup>, M. Giovannini<sup>17</sup>, M. Hocke<sup>18</sup>, A. Ignee<sup>19</sup>, E. M. Jung<sup>20</sup>, A. S. Klauser<sup>21</sup>, N. Lassau<sup>22</sup>, E. Leen<sup>23</sup>, G. Mathis<sup>24</sup>, A. Saftoiu<sup>25</sup>, G. Seidel<sup>26</sup>, P. S. Sidhu<sup>27</sup>, G. ter Haar<sup>28</sup>, D. Timmerman<sup>29</sup>, H. P. Weskott<sup>30</sup>

### Affiliations

Affiliation addresses are listed at the end of the article.

### Bibliography

DOI <http://dx.doi.org/10.1055/s-0031-1281676>

Published online 2011  
Ultraschall in Med © Georg Thieme Verlag KG Stuttgart · New York · ISSN 0172-4614

### Correspondence

### Thematic sections



	Thematic Section	Chairperson
1	Introduction	F. Piscaglia – C. Nolsøe
2	Generalities	D. Cosgrove
3	Equipment	H. P. Weskott
4	Investigator's training	O. H. Gilja

### List of Abbreviations



AAA = Abdominal Aortic Aneurysm  
AUC = Area Under the Curve  
CE = Contrast Enhanced  
CECT = Contrast Enhanced Computed Tomography  
CEMRI = Contrast Enhanced Magnetic Resonance Imaging

Unauthorized distribution is strictly prohibited.

Ultraschall Med / EJU Aug. 2011




# Indications - Liver

- Characterization of benign FLLs
  - FNH, hemangiomas, adenomas
- Detection of focal lesions
  - Metastasis
- Study FLL in cirrhosis
- Guiding of biopsies
- Guiding of intervention,- e.g. ablation



# 3 (4) Phases in liver perfusion

- Arterial phase
  - 0-30 sec.
- Portal phase
  - 30-120 sec.
- Sinusoidal phase
  - 2-4 min
- Post-vascular phase
  - 4-30 min






# Focal Liver Lesions

	Type of lesion	Arterial phase	Portal phase	Sinusoidal phase (parenchimal)
<b>Benign</b>	Haemangioma	Globular enhancement from the periphery	Centripetal filling	Progressive enhancement (iso to hyperechoic)
	Focal Nodular Hyperplasia	1. Strongly hyperechoic 2. In 40% of cases spoke and wheel pattern	Moderately hyperechoic or Isoechoic	Moderately hyperechoic or Isoechoic (central scar visible in 40% of cases)
	Adenoma	Strong homogeneous enhancement of short duration (capsular vessels)	Isoechoic	Isoechoic
<b>Malignant</b>	Hepato-cellular Carcinoma	Enhancement 1.Homogeneous 2.Inhomogeneous	Slightly hypoechoic	Slightly or strongly hypoechoic
	Hypervascular Metastases	1. Hyperechoic 2. Possible central area of necrosis in large lesions	Slightly hypoechoic	Strongly hypoechoic
	Hypovascular Metastases	1.No enhancement 2.Peripheral rim	Slightly hypoechoic	Strongly hypoechoic




# Lesion in Liver – S7






# Hyper-echoic Tumor






# Typical location: By the capsule or by an hepatic vein





# Referred from the CT-Lab Haemangioma ?






# Peripheral Globular Enhancement



...with slow centripetal filling




# Focal Nodular Hyperplasia - FNH



- FNH- a centrifugal stellate branching in early arterial phase
- Spoke wheel pattern in approx 40%
- Intense homogenous uptake
- Iso- or hyperechoic lesion is seen in portal venous phase.
- With these characteristic features:
  - sensitivity and specificity of contrast-enhanced low MI real-time US are 87.6% and 94.5%, respectively
  - Di Stasi 1996



# CEUS – Sonazoid - FNH






# Arterial Phase: 12 sec.






# Arterial Phase: 13 sec





# Arterial Phase: 13 sec






# Arterial Phase: 15 sec






# Liver cell adenoma



- Liver cell adenoma (LCA) is a rare primary benign neoplasm found mainly in young women with a history of oral contraceptive use
- The hypervascularity of adenomas can be demonstrated on Doppler,- sentripetal
- CEUS identification of the early and homogeneous hyperechoic enhancement in the periphery of the tumor, reflecting the presence of the subcapsular feeding arteries.
- The enhancement of LCA in the portal and late phases is nearly comparable with that of liver parenchyma, but LCA can remain slightly hypoechoic in relation to the adjacent liver





# CEUS - Real-time Perfusion



Dynamic abilities outperforms CT and MR




# A common problem: Focal Lesions in Fatty Livers





# CEUS in FLL




T1: 3:18



# Ultrasound in HCC

- Diagnosis
  - B-mode
  - Doppler
  - CEUS
- US-guided biopsy
- Per-operative guiding of ablation
- Follow-up and monitoring of treatment
- Surveillance / Screening






# HCC facts

## HCC – The great imitator

- Increasing incidence world-wide
- 90% has known ethiology
- AFP has limited sensitivity (approx. 60%)
- Most frequent: Alcohol (25%) and HCV
- NASH is increasing in incidence, thus feeding the HCC growth
- Barcelona criteria: 2 independant imaging methods are needed to avoid biopsy




# A patient with cirrhosis






# «The Basket Sign»






# CEUS of tumor

01/03/19 11:00:03 ADM C1-6 Sonozoid MI 0.30 TIs 0.0





# Any wash-out in late phase ?





# FNH versus HCC




**FNH**



**HCC**



# Early wash-out typical for mets




Not easily observed on CT and MRI



# Liver Metastasis ?

Before contrast injection






# Liver Metastasis ?

After contrast injection in late phase






# Mets from Rectal Cancer





# High-Frequency 9 MHz LA probe Post-vascular phase (Kupffer)




T1: 11:02



# One stop shopping

- US B-mode
- Doppler
  - Color flow
  - Pulsed Doppler
- Elastography
  - Shear wave
  - Strain imaging
- CEUS
- US-guided biopsy





# "Yes, we scan"





# US first = Ultrasound first...



”This is NOT fake news !”